IF YOU RECEIVED THIS FAX IN ERROR, PLEASE CALL 604-806-8886 IMMEDIATELY



PRESCRIBER'S ORDERS

NO DRUG WILL BE DISPENSED OR ADMINISTERED WITHOUT A COMPLETED **CAUTION SHEET**

DATE ND TIME	OPHTHALMOLOGY OPERATIVE ORDERS [see corresponding Medication Administration Record PHC-PH270-MA (R. Jan 17-18)] (Items with check boxes must be selected to be ordered)			
		OPERATIVE EYE:	☐ Left ☐ Right	
	PRE-OPERATIVE			
	Monitoring: Temp, BP, Pulse, RR, O ₂ Saturation. Monitoring as per DT Procedure Rooms, Pa Pre-operative Assessment protocol			atient Admission and
	Capillary Blood Glucose on admission (Diabetic patient)			
	MEDICATIONS:			
	Anxiety:	lorazepam ☐ 0.5 mg ★OR★ ☐ 1 mg ☐ lorazepam 0.5 to 1 mg sublingually PRN c	• •	
	Ophthalmic:	 □ tetracaine 0.5% ophthalmic solution x 1 drop into operative eye once □ cyclopentolate 1% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes □ moxifloxacin 0.5% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes □ PHENYLephrine 2.5% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes □ If slow dilation: PHENYLephrine 10% ophthalmic solution x 1 drop into operative eye once □ atropine 1% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes □ pilocarpine 2% ophthalmic solution x 1 drop into operative eye; repeat dose after 5 to 15 minutes □ lidocaine 2% jelly to be applied to operative eye by surgeon 		
	PROCEDURAL SEDATION:			
		Saline lock	moa	
	☐ fentanyl 25 mcg IV PRN to maximum of 50 mcg ☐ midazolam 0.5 mg IV PRN to maximum of 1 mg			
	Monitor patient as per Procedural Sedation in Clinics and Procedure Rooms, protocol			
		Discharge patients when procedural sedation Procedure Rooms, protocol	•	
	RETRO OR PERIBULBAR BLOCKS: (administered by ophthalmologist)			
		Saline lock		
		lidocaine 2% with epinephrine 1:100,000	¹to 4 mL *AND*	
		bupivicaine 0.75% 2 to 4 mL; may repeat to maximum of 8 mL co	nbined total	
	POST-OPERATIVE:			
	Monitoring: Monitor patient as per DT Procedure Rooms, Post op care discharge scoring assessment			
	acetazolamide 250 to 500 mg PO once immediately post-op			
	acetaminophen 325 to 650 mg PO Q3 to 4H PRN pain			
	☐ TYLENOL #3 1 to 2 tablets PO Q4H PRN pain			
	patient must remain supine x minutes post-operatively			
	Printed Name	Signature	College ID	Contact Number